

Autocorrelation functions in 3D Fully Frustrated Systems

G. Franzese^a, A. Fierro^a, A. De Candia^a and A. Coniglio^{a,b}

Dipartimento di Scienze Fisiche, Università di Napoli, Mostra d'Oltremare Pad.19 I-80125

Napoli, Italy

^aINFM - unità di Napoli and ^bINFN - sezione di Napoli

Abstract

We present a numerical study of autocorrelation functions of a 3D Fully Frustrated Ising model (FFIM) simulated by spin-flip Monte Carlo dynamics finding simple exponential decay for all the temperature above the critical temperature T_c for the autocorrelation of squared magnetization and stretched exponential decay for the energy autocorrelation below a temperature T^* with $T_c < T^* \leq T_p$ where T_p is the Kasteleyn-Fortuin and Coniglio-Klein percolation temperature. The results are compared to those on 2D FFIM to make light on the relevant mechanism in the onset of stretched exponential relaxation functions.

Autocorrelation functions of the form

$$f(t) = f_0 \exp -(t/\tau)^\beta \quad (1)$$

called stretched exponential or Kohlrausch-Williams-Watts (KWW) function are observed in glassy systems and spin glass (SG) well above the respective transition temperature and the origin of this kind of behaviour is still matter of debate.

For example in SG Randeria *et al.* [1] suggested that the onset of the non exponential relaxation function coincides with the critical temperature of the ferromagnetic Ising model and base their conjecture on the presence of non frustrated ferromagnetic-type clusters of interactions.

Furthermore Huse and Fisher [2] on the basis of droplet model and more recently Stauffer [3] on the basis of percolation arguments have predicted a KWW autocorrelation function also for ferromagnetic Ising model (FIM) in 2D for $T \leq T_c$ where T_c is the critical temperature. These arguments show that no such behaviour is expected in more than two dimensions and Stauffer has corroborated his prediction with simulations in 2 and 3D.

It is useful to note at this point that in FIM the critical temperature T_c coincides with the percolation temperature T_p of the Kasteleyn-Fortuin and Coniglio-Klein (KF-CK) clusters. [4]

These clusters can be obtained introducing bond with probability $p = 1 - e^{-2J/(k_B T)}$ between nearest-neighbor pair of spins satisfying the interaction (J is the interaction strength, k_B the Boltzmann constant and T the temperature) and represent correlated spins regions.

It would be interesting to investigate whether such prediction also holds for frustrated systems where the percolation temperature and the critical temperature do not coincide.

In particular in this work we considered the fully frustrated Ising model (FFIM). Here we resume the numerical results on 2D [5] and present those in 3D. [6]

The FFIM is a model with no disorder (unlike the SG and like the FIM) and with the interactions assigned in a regular way to give rise to *frustration*, *i.e.* a competition between them (like the SG and unlike the FIM). [7]

The FFIM Hamiltonian is given by

$$H = -J \sum_{\langle ij \rangle} (\epsilon_{ij} S_i S_j - 1), \quad (2)$$

where $\epsilon_{ij} = \pm 1$ are quenched regular assigned variables that give rise to frustration and $S_i = \pm 1$ are the Ising spins. [8] In this model the KF-CK clusters are such that they do not include a *frustrated loop*, that is a closed path of bonds which contains an odd number of interactions with $\epsilon_{ij} = -1$. These clusters overlap regions of not frustrated interactions and their percolation temperature is greater than the critical temperature. In Ref. [8] it is studied a general Hamiltonian, with Ising spins and s -state Potts spins, that has the FFIM as a particular case for $s = 1$. It is possible to see that for all $s \neq 1$ T_p of KF-CK clusters

corresponds to a thermodynamic transition of a s -state Potts model and that this thermal transition disappears for $s = 1$, *i.e.* for FFMI. This suggests that the KF-CK clusters may have an important role also for FFIM.

We simulate this model using the standard spin flip [9] dynamics.

In the 2D FFIM where $T_p > T_c = 0$ recent results [5] shows that stretched exponential autocorrelation functions are observed for $T \leq T_p$. Therefore the same qualitatively behaviour of 2D FIM is observed but in the FFIM is clear that the onset of stretched exponentials coincides, within numerical precision, with $T_p \neq T_c$, while in FIM is $T_p = T_c$.

To put more light on the problem we have done simulations on 3D FFIM.

To this goal we have calculated the autocorrelation function of squared magnetization M^2 and of energy E , where for a generic observable A the relaxation function is defined as

$$f_A(t) = \frac{\langle \delta A(t) \delta A(0) \rangle}{\langle (\delta A)^2 \rangle}, \quad (3)$$

where $\delta A(t) = A(t) - \langle A \rangle$, t is the time and $\langle \cdot \rangle$ is the thermal average.

We fit our data with a general function

$$f(t) = f_0 \frac{\exp -(t/\tau)^\beta}{t^x} \quad (4)$$

where f_0 , τ , β and x are parameters.

In Fig.1 we summarize the results for systems with linear size $L = 20$ and for temperatures below the estimated [10] $T_p(L)$ of KF-CK clusters in cubic FFIM.

For f_{M^2} the estimated parameters are $\beta \simeq 1$ and $x \simeq 0$ for all simulated temperatures below $T_p(L)$. For f_E there is a temperature T^* , between $T_p(L)$ and $T_c \simeq 1.355J/k_B$ (the estimated critical temperature [11]), below which is $\beta < 1$ and x weakly changes. Simulations on $L = 30 \div 50$ confirm these data.

Therefore the 3D FFIM results show a simple exponential decay in M^2 correlation functions but a stretched exponential decay in E correlation functions for $T \leq T^*$ with $T_c < T^* \leq T_p$. In conclusion the results on f_E in 3D FFIM, even if less clear than those on 2D FFIM, confirm the existence of an onset T^* of stretched exponential for the relaxation function well above the critical temperature T_c and almost equal (or slightly below) to

the KF-CK percolation temperature T_p that, in a generalized model with also s -state Potts variables with $s \neq 1$, corresponds to a thermodynamic transition of the Potts variables. [8]

REFERENCES

- [1] M.Randeria, J.P.Setha, and R.F.Palmer, *Phys. Rev. Lett.* **54**, 1321 (1985).
- [2] D.A.Huse and D.S.Fisher, *Phys. Rev. B* **35**, 6841 (1987).
- [3] D.Stauffer, *Physica A* **186**, 197 (1992) and P.Grassberger and D.Stauffer, *Physica A* **232**, 171 (1996).
- [4] C.M.Fortuin and P.W.Kasteleyn, *Physica* **57**, 536 (1972); A.Coniglio and W.Klein, *J.Phys.A* **13**,2775 (1980); A.Coniglio, F.Di Liberto, G.Monroy, and F.Peruggi, *Phys. Rev. B* **44**, 12605 (1991).
- [5] A.Fierro, A.De Candia and A.Coniglio, *Phys. Rev. E*, in press.
- [6] G.Franzese, A.Fierro, A.De Candia and A.Coniglio, in preparation.
- [7] D.C.Mattis *Phys.Lett.A* **56** 421 (1976); J.Villain *J.Phys.C* **10** 1717 (1977).
- [8] U.Pezzella and A.Coniglio, *Physica A*, **237**, 353 (1997); F.di Liberto and F.Peruggi to appear in *J. Phys. A*; G.Franzese and A.Coniglio, in preparation.
- [9] K. Binder, ed., *Monte Carlo Methods in Statistical Physics*, 2nd ed., (Springer, Berlin, 1986).
- [10] G.Franzese, in preparation
- [11] H.T.Diep, P.Lallemand and O.Nagai, *J.Phys.C*, **18**,1067 (1985).

FIGURES




FIG. 1. Correlation functions for $L = 20$: (a) f_{M^2} and (b) f_E at (from left to right) $k_B T/J = 3.5, 2.5, 1.9, 1.8$ from respectively 2,7,8,7 independent runs with statistics of $5 \cdot 10^6$ L^3 equilibrium update after $5 \cdot 10^3$ L^3 update of thermalization; insets: the corresponding estimated β and x in eq.(4) vs. $T/T_p(L)$ where $k_B T_p(L = 20)/J \simeq 3.95$. [10]